

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 3, 1997	3. REPORT TYPE AND DATES COVERED Final Technical, 11/15/94 - 12/31/96
4. TITLE AND SUBTITLE Fast Wavelet Transforms and the Numerical Solution of Initial Value Problems			5. FUNDING NUMBERS G N00014-95-1-0272
6. AUTHOR(S) Stanley J. Osher			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, Los Angeles Department of Mathematics 405 Hilgard Avenue Los Angeles, CA 90095			8. PERFORMING ORGANIZATION REPORT NUMBER 951602-01-A03
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research ONR 251 Ballston Centre Tower One 800 North Quincy Street Arlington, VA 22217-5660			10. SPONSORING / MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT

APPROVED FOR PUBLIC RELEASE. DISTRIBUTION IS UNLIMITED.

19970409 072

13. ABSTRACT (Maximum 200 words)

Fast algorithms to evaluate the approximate solution of time dependent problems were developed by taking advantage of the sparse wavelet representation of finite difference operators and using only part of the representation to compute the local solution. For example, we can evaluate the solutions at a point to parabolic equations with variable coefficients in $O(\log^4 N)$ operations when the equation has time independent coefficients. For time dependent coefficients; the complexity is $O(N \log^3 N)$. Additionally, high resolution numerical methods for the high frequency asymptotic expansion to electromagnetic propagation and scattering codes were developed. This replaces ray tracing by a direct solution to the eikonal equation. Moreover, we developed and solved generalized eikonal equations for diffraction phenomena.

DTIC QUALITY INSPECTED 2

14. SUBJECT TERMS Fast algorithms, sparse wavelet representations of finite difference operations, eikonal equations			15. NUMBER OF PAGES 1
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT unclassified	20. LIMITATION OF ABSTRACT unlimited

Final Report
ONR # N00014-95-1-0272

We developed fast algorithms to evaluate the solution of time dependent numerical methods for time dependent partial differential equations (PDEs) using multiscale analysis.

In [1] we developed very fast algorithms to evaluate the approximate solution to parabolic equations at one point without calculating in the whole space time domain. This was achieved by taking advantage of the sparse wavelet representation of finite difference approximations (which are analogous to those of Green's function for the operator) and using only parts of the representation to compute the local solution. The complexity for solving at a given point is only $O(\log^4 N)$ when the equation has time independent solutions. When the coefficients do depend on time, the complexity is $O(N \log^3 N)$.

Earlier, [2], we developed an algorithm which gives the solution with $O(N^d \log(N))$ complexity for linear hyperbolic equations in one space dimension and linear parabolic equation in d dimensions.

In a parallel effort we developed high resolution numerical methods for the high frequency asymptotic expansions to electromagnetic propagation and scattering [3]. This replaces ray tracing by a direct solution to the eikonal equation using new numerical methods for Hamilton-Jacobi equations and conservation laws developed here.

Additionally we developed and solved generalized eikonal equations for diffraction phenomena [4]. We were able to correct for boundary effects using a new perturbed geometric optics system.

Finally, we began a program in which the eikonal equation was generalized to accommodate solutions containing many phases. New equations were developed which are based on the same high frequency approximation of the scalar wave equation as the eikonal and transport equations.

References

- [1] A. Jiang and S. Osher, "Fast Wavelet Based Methods for Local Solutions to Parabolic Problems", SIAM J. Scientific Comp., (to appear).
- [2] B. Engquist, S. Osher, and S. Zhong, "Fast Wavelet Algorithms for Linear Evolution Equations", SIAM J. on Scientific Comput., v. 15, (1994), pp. 755-775.
- [3] E. Fatemi, B. Engquist, and S. Osher, J. Comput. Phys., v. 20, (1995), pp. 145-155.
- [4] E. Fatemi, B. Engquist and S. Osher, UCLA CAM Report #95-11, 1995.
- [5] B. Engquist and O. Runborg, J. Comput. and Appl. Math., v. 74, (1996), p. 175.